

Brazilian Primes

Jon Grantham
Hester Graves

Institute for Defense Analyses
Center for Computing Sciences
Bowie, Maryland

April 2019

Primes of the form $x^2 + x + 1$

- ▶ Last year at SERMON, I talked about computations of primes of the form $x^2 + 1$ for $x < 2.5 \times 10^{14}$.
- ▶ Previously, computed for $x < 10^{12.5}$ by Gerbicz and Wolf.
- ▶ This led to questions about other prime values of cyclotomic polynomials.

Primes of the form $x^2 + x + 1$

- ▶ Last year at SERMON, I talked about computations of primes of the form $x^2 + 1$ for $x < 2.5 \times 10^{14}$.
- ▶ Previously, computed for $x < 10^{12.5}$ by Gerbicz and Wolf.
- ▶ This led to questions about other prime values of cyclotomic polynomials.
- ▶ Let's ignore the cases $k = 1$ and $k = 2$ for now.

Primes of the form $x^2 + x + 1$

- ▶ Last year at SERMON, I talked about computations of primes of the form $x^2 + 1$ for $x < 2.5 \times 10^{14}$.
- ▶ Previously, computed for $x < 10^{12.5}$ by Gerbicz and Wolf.
- ▶ This led to questions about other prime values of cyclotomic polynomials.
- ▶ Let's ignore the cases $k = 1$ and $k = 2$ for now.
- ▶ For $k = 3$, the previous published computation appears to be up to 1.21×10^9 , by Poletti (1929).
- ▶ It was easy to modify the $x^2 + 1$ code to compute a table up to 10^{12} .
- ▶ Fun fact: $\phi_3(x - 1) = \phi_6(x)$, so we have done the $k = 6$ case.

An Interlude about Luigi Poletti



- ▶ Luigi Poletti (1864-1967) was an banker from Pontremoli in Italy who stumbled across a book of Derrick Lehmer at age 47.
- ▶ He spoke at the 1928 ICM.
- ▶ After World War II, he served on a commission to rebuild French science.
- ▶ He wrote original poems in and translated Dante into his native dialect (Pontremolese).
- ▶ There is a Via Luigi Poletti in Pontremoli.

An Interlude about Luigi Poletti



- ▶ Luigi Poletti (1864-1967) was an banker from Pontremoli in Italy who stumbled across a book of Derrick Lehmer at age 47.
- ▶ He spoke at the 1928 ICM.
- ▶ After World War II, he served on a commission to rebuild French science.
- ▶ He wrote original poems in and translated Dante into his native dialect (Pontremolese).
- ▶ There is a Via Luigi Poletti in Pontremoli.
- ▶ We are going to call primes of the form $x^2 + x + 1$ “Poletti primes”.

An Interlude about My Co-Author



Brazilian Primes

- ▶ Brazilian Primes are primes that are all 1s (repunits) in some base $b > 1$ (of length q at least 3).
- ▶ For primes $q > 2$, primes represented by the q th cyclotomic polynomial are Brazilian primes.
- ▶ Originated at the 1994 Iberoamerican Mathematical Olympiad in Fonseca, Brazil, in a problem proposed by the Mexican math team.
- ▶ First studied by Schott (2010).
- ▶ Thanks to Hester for translating from the French.

$$k = 5$$

- ▶ It is possible to modify the existing algorithm to compute primes of the form $x^4 + x^3 + x^2 + x + 1$.
- ▶ But to compute up to $x < B$, we need to sieve up to B^2 .
- ▶ So the running time is now $O(B^2 \log B \log \log B)$.
- ▶ In other words, a table for $B < 10^6$ would take as long as our $k = 3$ table for $B < 10^{12}$.

A better algorithm

- ▶ If we sieve up to B , we get numbers of the form $x^4 + x^3 + x^2 + x + 1$ which are B -rough.
- ▶ Heuristically, there should be $O(x/\log x)$ of these. (Buchstab)
- ▶ Need a fast way to distinguish primes from composites.

A better algorithm

- ▶ If we sieve up to B , we get numbers of the form $x^4 + x^3 + x^2 + x + 1$ which are B -rough.
- ▶ Heuristically, there should be $O(x/\log x)$ of these. (Buchstab)
- ▶ Need a fast way to distinguish primes from composites.
- ▶ The Pocklington–Lehmer test on N runs in $O(\log^2 N)$ time if you can fully factor a piece of $N - 1$ of size $N^{1/2}$.
- ▶ Here, $N = x^4 + x^3 + x^2 + x + 1$, so $N - 1 = x^4 + x^3 + x^2 + x = x(x + 1)(x^2 + 1)$. So you can!
- ▶ We can still generate a list up to B in time $O(B \log B \log \log B)$
- ▶ So verified up to 10^{12} .

A Conjecture

- ▶ Schott conjectured that there are no Brazilian Sophie Germain primes.
- ▶ Recall that a Sophie Germain prime is a prime p such that $2p + 1$ is also prime.
- ▶ If p is a Sophie Germain prime, then we say that $2p + 1$ is a “safe” prime.
- ▶ It is straightforward to show that if p is a Brazilian prime, then q is an odd prime.

A Lemma

- ▶ If p is a **Brazilian Sophie Germain prime**, $p \equiv q \equiv 2 \pmod{3}$ and $b \equiv 1 \pmod{3}$.
- ▶ If p is a Sophie Germain prime, then 3 cannot divide the safe prime $2p + 1$, so p cannot be congruent to 1 (mod 3).
- ▶ The number 3 is not Brazilian, so $p \neq 3$ and thus $p \equiv 2 \pmod{3}$.
- ▶ If $3|b$, then $p = b^{q-1} + b^{q-2} + \cdots + b + 1 \equiv 1 \pmod{3}$, which is a contradiction.
- ▶ q is an odd prime, so if $b \equiv 2 \pmod{3}$, then $p \equiv 1 \pmod{3}$, a contradiction.
- ▶ We conclude that $b \equiv 1 \pmod{3}$, so that $q \equiv p \pmod{3}$, and therefore $q \equiv 2 \pmod{3}$.

Finding Counterexamples

- ▶ So the key to looking for counterexamples is to look in our $k = 5$ list, not our list of Poletti primes.
- ▶ We find $28792661 = 73^4 + 73^3 + 73^2 + 73 + 1$ as the smallest example, and 104,890,302 examples up to 10^{46} .

Finding Counterexamples

- ▶ So the key to looking for counterexamples is to look in our $k = 5$ list, not our list of Poletti primes.
- ▶ We find $28792661 = 73^4 + 73^3 + 73^2 + 73 + 1$ as the smallest example, and 104,890,302 examples up to 10^{46} .
- ▶ (There are 104,890,282 examples up to 10^{46} .)
- ▶ There are only 20 other Brazilian Sophie Germain primes up to 10^{46} , all of length 11.

Finding Counterexamples

- ▶ So the key to looking for counterexamples is to look in our $k = 5$ list, not our list of Poletti primes.
- ▶ We find $28792661 = 73^4 + 73^3 + 73^2 + 73 + 1$ as the smallest example, and 104,890,302 examples up to 10^{46} .
- ▶ (There are 104,890,282 examples up to 10^{46} .)
- ▶ There are only 20 other Brazilian Sophie Germain primes up to 10^{46} , all of length 11.
- ▶ The first few counterexamples were discovered independently by Giovanni Resta and Michel Marcus.

Finding Counterexamples

- ▶ So the key to looking for counterexamples is to look in our $k = 5$ list, not our list of Poletti primes.
- ▶ We find $28792661 = 73^4 + 73^3 + 73^2 + 73 + 1$ as the smallest example, and 104,890,302 examples up to 10^{46} .
- ▶ (There are 104,890,282 examples up to 10^{46} .)
- ▶ There are only 20 other Brazilian Sophie Germain primes up to 10^{46} , all of length 11.
- ▶ The first few counterexamples were discovered independently by Giovanni Resta and Michel Marcus.
- ▶ See A306845 in the On-Line Encyclopedia of Integer Sequences.

Conditional Results

- ▶ Recall that Schinzel's Hypothesis H says that any set of polynomials, whose product is not identically zero modulo any prime, is simultaneously prime infinitely often.
- ▶ Assuming Hypothesis H, there are infinitely many Brazilian Sophie Germain primes.

Conditional Results

- ▶ Recall that Schinzel's Hypothesis H says that any set of polynomials, whose product is not identically zero modulo any prime, is simultaneously prime infinitely often.
- ▶ Assuming Hypothesis H, there are infinitely many Brazilian Sophie Germain primes.
- ▶ Assuming the Bateman–Horn conjecture, the number of Brazilian Sophie Germain primes is $\sim C \frac{x^{1/4}}{\log x^2}$, for some C .
- ▶ (Look at the $k = 5$ case, and show that it dominates the others.)

A Related Proposition

- ▶ **Proposition:** The only Brazilian prime which is a safe prime is 7.
- ▶ If $p = b^{q-1} + \dots + b + 1$ is a safe prime, then $\frac{p-1}{2} = \frac{1}{2}(b^{q-1} + \dots + b)$ must also be prime.
- ▶ This expression, however is divisible by $\frac{b(b+1)}{2}$, which is only prime when $b = 2$ and $p = 7$.

Future Work

- ▶ Extend tables?
- ▶ $k = 7$ requires use of Brillhart–Lehmer–Selfridge test (factorization of $N - 1$ up to $N^{1/3}$).
- ▶ Find application of Konyagin–Pomerance (which works with $N^{3/10}$).